Monoclonal antibody therapeutics: Leading companies to maximise sales and market share

Karl Ziegelbauer and David R. Light
Date Received (in revised form): 26th October, 2007

Karl Ziegelbauer

is a Vice President for Protein Therapeutics at Bayer HealthCare AG. He received his PhD from the University of Tübingen. For the past 14 years, he has been involved in drug discovery in various positions with increasing responsibilities in Germany, Japan, and the United States. His experience covers all aspects of drug discovery in various indications up to proof of concept in humans as well as lifecycle management and research alliance management.

David R. Light

received his PhD from the University of Wisconsin and completed a fellowship at the Massachusetts Institute of Technology before joining the newly formed Biocatalysis Department (later Genencor) in Genentech. He joined Berlex, a wholly owned subsidiary of Schering AG, as a protein chemist to develop biologic therapeutics including soluble thrombomodulin in Cardiovascular Research and later was project team leader of the factor Xa inhibitor project. Most recently, he headed the Antibody Technology Group focussing on antibody targets in cancer before joining the Protein Therapeutics group in Bayer.

Abstract

A close look at the biology and pharmacology of monoclonal antibodies reveals both their continuing promise as therapeutic agents to address unmet medical needs, as well as a number of challenges to the future discovery and development of this unique class of biologics. A remarkably consistent experience of reliable clinical efficacy and safety ensures that Biotech and Pharma have strong incentives to accelerate the antibody drug discovery process. Their attractive commercial potential invites consideration of potential challenges to the future expansion of the monoclonal antibody drug market. Four challenges arise from scientific and technical aspects of the antibody drug format: drug target limitations, biodistribution limitations, species specificity issues, and limitations to the route of administration and four challenges are based in the commercial and clinical use of antibody drugs: cost of goods, product differentiation within the antibody market, competition from small molecule drugs, and price sensitivity of clinical acceptance. Despite these challenges and recent setbacks, such as the withdrawal and subsequent relaunch of Tysabri and the TGN1412 Phase I disaster, the prevailing opinion is that monoclonal antibodies will continue to be safe and effective medicines that are worthy of commercialisation.

Journal of Commercial Biotechnology (2008) 14, 65–72. doi:10.1057/palgrave.jcb.3050081; Published online 27 November 2007;

Keywords: monoclonal antibody drugs, biologic therapeutics, antibody market

Correspondence: Karl Ziegelbauer, Protein Therapeutics, Bayer HealthCare AG, Wuppertal 42096, Germany

Tel: +49 202 36 5293 Fax: +49 202 36 4585

E-mail: karl.ziegelbauer@bayerhealthcare.com

INTRODUCTION

The usage of immune horse antiserum to treat severe infectious diseases such as tetanus and diphtheria at the end of the 19th century was



the first application of therapeutic antibody drugs. Over a century of experience with biological drug products derived from human or animal serum followed, including polyclonal immunoglobulin antibodies to treat severe infections and snake bites, as well as the development of serum protein drug products to treat genetic or acquired deficiency diseases such as diabetes and haemophilia. This valuable knowledge base in the production, formulation, delivery, and clinical usage of serum-derived protein drugs set the stage for the rapid expansion of biotechnology-derived protein therapeutics beginning in the 1980s driven by advances in both recombinant DNA and monoclonal antibody technologies. These early years also previewed challenges that continue to impact the development of protein-based therapeutics including the formation of antibodies by patients that may neutralise the activity of a biological drug or trigger adverse reactions ranging from mild fever to life-threatening anaphylactic shock during repeated treatments.

Over two decades of the current biotechnology era has been characterised by intense research to develop antibody-based drugs beginning with the first clinical testing of mouse monoclonal antibodies. Now the industry standard has shifted away from nonhuman and chimaeric antibodies to focus on fully human or humanised antibody drug candidates. Significant advances in the discovery of human monoclonal antibody drug candidates have resulted in novel approaches utilising both in vivo and in vitro methods. In vivo approaches focus the power of the vertebrate adaptive immune system to directly create potent human antibodies in transgenic animals or non-human antibodies that are subsequently humanised. In vitro approaches are driven by advances in antibody library design¹ and these antibody libraries are subsequently used for selecting and increasing the affinity of human antibodies displayed on ribosomes, phage, bacteria, yeast, and mammalian cells derived from human B-cells. The potential of this accumulated technology

is truly impressive, allowing the formation of polyclonal mixtures of fully human monoclonal antibodies with a thousand-fold improved potency and the potential to treat severe infectious diseases such as botulism.²

THE CURRENT ANTIBODY DRUG MARKET

The natural role of antibodies is to block infectious disease by binding foreign agents or infected cells leading to activation of the host immune system. Only one of the 21 current marketed drugs (Synagis, respiratory syncytial virus, MedImmune) and none of 38 advanced clinical antibody drug candidates, however, are designed for this type of application.³ Some antibody drugs or drug candidates that target leukaemia, lymphoma, or severe autoimmune disease do bind to B-cell or T-cell-associated surface antigens (CD-20 or CD-52) in order to trigger depletion of their target cells by complement or Fcyreceptor-mediated mechanisms; however, cell depletion by the immune system is not the goal of the majority of antibody therapeutics. Instead, the field has advanced by discoveries that create new functionalities for monoclonal antibodies, including receptor binding to modify activation, prevent dimerisation, trigger internalisation, block proliferation, or induce apoptosis, binding, and neutralisation of cytokines or growth factors, and targeting chemical, protein, or radioactive toxins to target cells. Largely as a result of the discovery of these and other novel functionalities, antibody drugs are now breakthrough therapies for a variety of diseases, especially in the area of oncology and severe immunological disease indications.

The current market size for monoclonal antibodies is estimated to be over \$20bn (US). This market is dominated by five antibody drugs Avastin (bevacizumab), Herceptin (trastuzumab), Humira (adalimumab), Remicade (infliximab), and Rituxan (rituximab), which together account for ~80 per cent of market. In some estimates, the

antibody market is projected to grow to \$30bn or even more over the next 3–6 years, driven mainly by oncology applications.^{4,5} This growth will include a mixture of new therapeutic applications for marketed antibody drugs, improved antibodies aimed at clinically validated targets, and the introduction of novel antibody drugs to novel targets.

Recent trends indicate that large Biotech increasingly relies on an ageing portfolio of approved protein therapeutics;6 therefore, these projected market increases may need to be adjusted if the rate of successful launches leading to novel antibody drugs begins to slow. Biotech and Pharma, however, have a compelling incentive to accelerate the antibody drug discovery process because the commercial aspects of this drug modality remain very impressive. Therapeutic antibodies have a high drug approval success rate once they reach clinical testing (29 per cent for chimaeric antibodies, 25 per cent for humanised antibodies compared to a success rate of approximately 11 per cent for small molecules).7 In addition, much of the development and clinical experience that is gained from the generation and optimisation of one antibody product can be readily applied to subsequent therapeutic antibodies, diminishing some of the development, manufacturing, and clinical risks that are intrinsic to drug development.

Owing to their exquisite specificity and ability to affect unique biological functions, monoclonal antibodies have the potential to provide a continued source of effective, safe, and reliable therapies. The introduction of such new therapies will benefit patients having a variety of debilitating diseases that otherwise respond poorly to alternate approaches. Based on the impact of the successful discovery of novel antibody functions on the current portfolio of antibody drugs, it is likely that the ability to continue to engineer novel functionalities by using new antibody formats will drive the expansion of the antibody drug market in the future.

POTENTIAL CHALLENGES TO THE FUTURE OF ANTIBODY DRUGS

The discovery of a continued stream of monoclonal antibody-based therapies offers tremendous opportunities for Pharma and Biotech companies, but also harbours a variety of scientific and commercial challenges. Currently, 21 monoclonal antibodies are approved for therapeutic use, 11 of which are humanised, five chimaeric, three of murine origin, while only two are fully human antibodies. In contrast, of 38 antibodies in advanced clinical testing, 31 are either humanised (14) or fully human (17). These numbers confirm that human antibodies are now the standard of the industry, whether obtained by in vivo (immunisation) or in vitro (antibody display) methods. Given the large number of antibodies in clinical trials or preclinical development, it is clear that technologies for the discovery of human antibodies are not rate limiting. Rather, the key to success will either be to identify the most effective, novel, and proprietary target in a complex pathological setting, or to identify a more effective approach to a known target, and both will be guided by emerging target validation approaches.

It is instructive to consider eight challenges to the future expansion of the monoclonal antibody drug market. Four of these challenges arise from scientific and technical aspects of the antibody drug format: drug target limitations, biodistribution limitations, species specificity issues, and limitations to the route of administration. The remaining four challenges arise from a consideration of the commercial and clinical usage of antibody drugs: cost of goods, product differentiation within the antibody market, competition from small molecule drugs, and price sensitivity of clinical acceptance.

Drug target limitations

Following intravenous injection, antibody drugs access targets in the extracellular and vascular

米

space. The Fc domain of the IgG format interacts with the endothelial FcRn receptor, facilitating access to the perivascular space after transient movement through the endothelium; however, the interior of these endothelial cells is not targeted. Antibody immunotoxin conjugates utilise antibody internalisation following receptor binding, but the goal of this application is delivery of the toxin payload to the intracellular space, not delivery of the antibody itself, which is rapidly hydrolysed. Disulphides that are needed to maintain the dimeric structure of the IgG format break down in the reducing environment of the intracellular cytosol and deactivate the potent binding of the antibody. Thus, antibody drugs are limited to extracellular targets.

The consequence of this limitation can be appreciated by considering the rich list of intracellular drug targets, several of which are targeted by small molecule drugs that are market leaders. These intracellular targets include HMG-CoA reductase (statins), nuclear hormone receptor agonists or antagonists (glucocorticoid, oestrogen, progesterone, etc), phosphodiesterase (PDE5), immunophilins (cyclosporine, FK506), and kinases (receptor tyrosine kinase, thymidine kinase). Even when limited to the extracellular arena, there appears to be a preference for prominent antigens on extracellular domains or soluble ligand targets. Thus, a notable omission from the list of extracellular targets that have been addressed by advanced antibody candidates include the seven-transmembrane receptors. In this large and ubiquitous family of membrane receptors are the G-protein coupled receptors (GPCR) that are often therapeutic targets for small molecule drugs. Technical advances may expand the range of extracellular targets for antibody drugs to include the GPCRs; however, it is not likely that the natural limitations of antibodies will be overcome to allow access to valuable intracellular targets.

Biodistribution limitations

Significant biodistribution and tissue penetration challenges limit the application of

antibody drugs. Solid tumours make up the majority of human cancers (~85 per cent). To date, nine antibodies have been approved for the treatment of human cancers, but only three target solid tumours and one of these, bevacizumab, is actually directed towards a soluble ligand target, not to its cell surface receptor expressed on cells inside the solid tumour tissue. This suggests that additional barriers are associated with the treatment of solid tumours that are not present for haematological malignancies. These limitations need to be addressed before the successful treatment of human solid tumours by antibody drugs can expand. Even greater challenges may exist for other disease indications in privileged tissue like the central nervous system. When faced with these challenging biodistribution applications, the high-molecular-weight IgG format may ultimately fail to achieve the required tissue penetration. In the future, this natural antibody format may be replaced by protein drugs derived from alternate antibody formats or antibody mimetic scaffolds. It remains to be determined whether these molecules are able to more effectively access poorly vascularised tumours or tissues protected by the blood-brain barrier.

Species specificity issues

One of the valuable properties of antibody drugs is their exquisite specificity, allowing them to bind one particular epitope in the presence of many other similar binding targets. This may, however, lead to extended preclinical development times for antibody drugs because antibodies to a human target may not bind the similar target molecule in species commonly used for efficacy or safety testing (mouse, rat, rabbit, dog, etc). In addition, well-known recent clinical results indicate that the species specificity of Fc receptor binding must be given greater consideration during development of future antibody drugs.

There is not one universally accepted solution to the issue of species specificity of

antibody epitope recognition; however, there are a number of potential solutions, all of which add time or require significantly more resources for antibody drug discovery. One potential solution is to develop two separate antibodies in parallel in order to include one antibody that recognises the antigen in a species used for animal testing. Another solution is to establish transgenic mice expressing both the human target protein and the necessary human auxiliary proteins in order to allow testing of human-specific antibodies in transgenic mouse models. A third potential solution is to screen antibodies that recognise both the human and the mouse antigen and advance only those candidate antibodies with dual species specificity. This option harbours the potential risk of discarding unique antibodies to a functional human epitope not found on the mouse homologue. A final potential option would be to complete the entire concept validation studies in vitro utilising human cell or tissuebased models. Then, the first in vivo proof of concept would be in human clinical studies; however, the acceptance of this approach is problematic in the post-TGN1412 era.

The major lesson arising from the disastrous clinical testing of TGN1412 (anti-CD28 IgG T cell superagonist) is that special caution is needed in the design and execution of 'first in man' clinical trials.8 In addition, however, based on this tragic event, an increased emphasis on preclinical consideration of Fc receptor interactions is likely to be required by regulatory agencies.9 The 'cytokine storm' was not found during preclinical testing of TGN1412 in cynomolgus monkeys, indicating that species differences in Fc receptor binding may be important for IgG tests even in nonhuman primates. Functional aspects of the traditional IgG format can be finely tuned by post-translational glycosylation in a proprietary production cell line or by Fc engineering to selectively trigger effector functions such as antibody-dependent cellular cytotoxicity. It is clear that the Fc portion of the monoclonal antibody plays a significant role in the

extended serum half-life that is now expected for this class of biological drugs. Thus, Fc modifications must achieve the right balance of clearance, Fc receptor functionality, and clinical safety. Modification or elimination of Fc receptor binding altogether may be included as a motivation to pursue alternatives to the traditional IgG format. Various new antibody formats as well as mimetics are being pursued, but there is limited or no clinical experience with most of these antibody formats and mimetics. The jury is still out on whether one of the many emerging novel scaffolds that eliminate Fc receptor binding can effectively substitute for monoclonal antibodies in the mid-term. It is likely, however, that preclinical development issues based on species-specific target recognition will continue to challenge both antibody and antibody mimetic-based drug candidates.

Limitations to the route of administration

Four of the top five selling antibody drugs and 16 of the 21 approved antibody drugs are administered by intravenous infusion. Several antibody drugs are approved for subcutaneous injection. Selected antibodies either are approved or have been successfully tested in the clinic using other routes of administration often designed for a specific indication including intramuscular (palivizumab), intravitreal (ranibizumab), intracoronary (abciximab), and intraperitoneal, intraventricular, or intralesional (rituximab). Subcutaneous injection offers the possibility of self-injection by the patient; however, a comparison of the TNFα antagonists infliximab (intravenous infusion) and adalimumab (subcutaneous injection) showed equivalent short-term efficacy despite this difference in the route of administration. 10 Over the course of long-term treatment, self-administration and ease of use may be anticipated to impact patient compliance and acceptance of antibody drugs. Insulin represents a protein therapeutic that is readily



delivered by subcutaneous injection, but over the years more acceptable delivery methods have been actively sought due to market demand. In the case of insulin, both the popular insulin pen technologies utilising extremely small, short needles as well as inhaled insulin powder (Exubera) have subsequently been developed. When the relatively low molecular weight of insulin (~5,800 Da) is compared to the high molecular weight of the IgG format (~150,000Da) found in the majority of approved antibody drugs, it is clear that further technical advances will be needed to achieve the ease of use of pen injection or inhalation technologies. An additional potential benefit of the lower molecular weight of either alternate antibody formats (~25,000-75,000 Da) or antibody mimetics $(\sim 9,000-15,000 \, \text{Da})$ may be to allow expanded routes of administration. If any of these improved routes of administration are realised they will represent an important milestone for the development of these alternatives to traditional antibody drugs.

Cost of goods (COGS)

Significant costs are associated with the identification, optimisation, and production of monoclonal antibodies due to the cost of manufacturing and intellectual property considerations. Because of their complex structure, monoclonal antibodies in the IgG format are generally limited to production in mammalian cells. These large protein drugs (~150,000 Da) require post-translational modifications and critical disulphide bonds for full activity. The usual route of production in either Chinese hamster ovary (CHO) or mouse myeloma (NS0) cell lines is often expensive and time consuming. The technologies to discover and produce monoclonal antibodies have been heavily patented and companies that are active in this field often need to acquire one or more licenses, either research or commercial. These intellectual property costs are not associated with a single patent and not limited to the

antibody molecule itself, but may include a collection of technologies needed for antibody drug generation, optimisation, and production. These technologies may include, among others, affinity maturation, humanisation methods, the expression systems (promoter and poly A sequence), and cell lines used to produce the antibody with the appropriate post-translational modifications needed to ensure the desired functionality. The cumulative costs associated with licensing these technologies are often referred to as 'stacking royalty' payments.

Product differentiation within the antibody market

Within the TNFa antagonist arena, different routes of administration have not yet distinguished similar antibody products with respect to short-term efficacy. This indication is not unique in having several approved antibody drugs or antibody drug candidates in advanced development. Marketed antibody drugs and advanced candidates are often directed to the same target and there are apparently four TNFa antagonists (infliximab, adalimumab, golimumab, certolizumab) in addition to the Fc fusion protein etanercept, five antibodies targeting the B cell receptor CD20 (rituximab, ibritumomab, tositumomab, ofatumumab, ocrelizumab), five to the EGF receptor EGFR (cetuximab, panitumumab, matuzumab, nimotuzumab, zalatumumab), and three to VEGF/VEGFR signalling (bevacizumab, ranibizumab, CDP-791) in addition to the Fc fusion VEGF-Trap. A simple list is not a fair comparison and in some cases the antibody formats are different (immunotoxin versus naked antibody, or IgG versus Fab), the indications are different (solid tumour versus macular degeneration, lymphoma versus arthritis), or the approach to the target is different (soluble ligand versus receptor extracellular domain); however, it is clear that the potential to differentiate between similar antibody products will represent a challenge to the industry.

Competition from small molecule drugs

In addition to the need to differentiate antibody drugs from each other, there is emerging pressure from small molecule drugs directed toward similar targets as the antibodies. No good solution has been found to the daunting technical challenge of finding small molecules that can bind and block tight protein-protein interactions. Antibodies are excellent agents for binding one protein and preventing binding of its biological partner. Small molecule drugs are not likely to compete directly with antibodies using this mechanism, but the therapeutic targets of antibody drugs may be addressed by alternate mechanisms taking advantage of the ability of small molecule drugs to antagonise intracellular targets. Growth factor receptors often are comprised of an extracellular domain suitable for antibody binding and an intracellular kinase domain that may be antagonised by small molecule inhibitors. Lapatinib (Glaxo Smith Kline) is an example of one such oral receptor tyrosine kinase inhibitor that targets both EGFR (ErbB-1) and HER2 (ErbB-2) receptors and that may compare favourably with antibody drugs to these targets. Similarly, by inhibiting the tyrosine kinase domains of VEGFR-1, VEGFR-2, and VEGFR-3 and other receptors (Raf, PDGFR-B, KIT, FLT-3, and RET), the oral drug Nexavar (Bayer) may compare favourably to antibodies targeting VEGF/VEGFR signalling. In the future, combined treatment that includes both antibody and small molecule drugs may prove most effective for life-threatening diseases. It is likely, however, that further challenges to the ability of antibodies to address drug targets will arise from advances in the ability to identify small molecule leads to the same targets.

Price sensitivity of clinical acceptance

The cost issues for antibody therapies do not end when the antibody drug has been successfully produced and packaged. Antibody drugs targeting cancer rarely cure this disease, especially in the advanced stages of cancer. For maximum benefit, antibody cancer drugs are usually administered in combination with chemotherapy or radiotherapy. These combined treatments significantly increase the total cost to the patient. For example, the FOLFOX regimen (Fluorouracil, leucovorin, and oxaliplatin) costs nearly \$12,000 dollars for an 8-week course compared to approximately \$21,000 for FOLFOX combined with the antibody drug bevacizumab, but the combination with antibody drug results in a significant increased benefit in the median survival time.11 Improved benefits in the clinic will drive Biotech and Pharma to work with clinicians to find ways to continue to improve the costs of antibody therapies. National healthcare providers, however, may be reluctant to pay for the high cost of antibody drugs. Recently, the National Institute for Health and Clinical Excellence in the United Kingdom was widely criticised when it failed to recommend cetuximab and bevacizumab for advanced bowel cancer based on their estimate that these drugs were not cost-effective in the treatment of metastatic colorectal cancer. 12 Thus, a final challenge will be to identify antibody drugs that are efficacious therapeutics and that are recognised as cost-effective by healthcare providers.

In summary, monoclonal antibodies, along with their derivatives and conjugates, provide tremendous opportunity in the mid-term for the discovery of new treatments for diseases with high unmet medical need. Despite recent setbacks such as the withdrawal and subsequent relaunch of Tysabri and the TGN1412 Phase I disaster, the prevailing opinion is that monoclonal antibodies will continue to be safe and effective medicines. Emerging technologies in novel antibody formats and mimetics will further provide opportunities to improve this unique class of biological drugs and will help to ensure their continued commercial success.



References and Notes

- Fuh, G. (2007). Synthetic antibodies as therapeutics. Expert. Opin. Biol. Ther. 7(1), 73-87.
- Garcia-Rodriguez, C., Levy, R., Arndt, J. W., Forsyth, C. M., Razai, A., Lou, J., Geren, I., Stevens, R. C. & Marks, J. D. (2007). Molecular evolution of antibody cross-reactivity for two subtypes of type A botulinum neurotoxin. *Nat. Biotechnol.* 25(1), 107–116.
- 3. While the application of the clinical candidate TNX-355 (Genentech/Tanox) is to block HIV infections, the epitope is not on the HIV virus itself, but on human CD4, a receptor that facilitates entry into target cells.
- Baker, M. (2005). Upping the ante on antibodies. Nat. Biotechnol. 23(9), 1065-1072.
- Wiles, M. & Andreassen, P. (2006). Monoclonals The billion dollar molecules of the future. *Drug. Disc. World* 7(4), 17–23.
- Jarvis, L. M. (2007). Biotechs rely on old products. Chem. Eng. News. 85(34), 34–35.
- 7. Carter, P. J. (2006). Potent antibody therapeutics by design. Nat. Rev. Immunol. 6(5), 343-357.
- 8. Duff, G. W., Lechler, R., Borysiewicz, L., Inglis, S., Parry, V., Walport, M., Markham, A. Haites, N.,

- McMichael, A., Hepple, B., Piromohamed, M., Loewer, J., Klareskog, L., Park, K. & Waldmann, H. (2006). Final report of the Expert Scientific Group (ESG) on phase one clinical trials. The Stationary Shop (www.tsoshop.co.uk).
- Stebbings, R., Findlay, L., Edwards, C., Eastwood, D., Bird, C., North, D., Mistry, Y., Dilger, P., Liefooghe, E., Cludts, I., Fox, B., Tarrant, G., Robinson, J., Meager, T., Dolman, C., Thorpe, S. J., Bristow, A., Wadhwa, M., Thorpe, R. & Poole, S. (2007). "Cytokine Storm" in the phase I trial of monoclonal antibody TGN1412: Better understanding the causes to improve preclinical testing of immunotherapeutics. J. Immunol. 179(5), 3325–3331.
- Schwartzman, S. & Morgan, G.J. (2004). Does route of administration affect the outcome of TNF antagonist therapy. Arthritis. Res. Ther. 6(Suppl 2), S19-S23.
- Schrag, D. (2004). The price tag on progress Chemotherapy for colorectal cancer. N. Engl. J. Med. 351(4), 317–319.
- Hawkes, N. (2006). Patients angered as watchdog refuses to allow bowel cancer drugs on NHS. The Times, 21 August.

FROM THE ANALYST'S COUCH

Monoclonal antibodies market

Janice Reichert and Alex Pavlou

Following the success of recombinant proteins, therapeutic monoclonal antibodies (mAbs) represent the second wave of innovation created by the biotechnology industry during the past twenty years. Between 2001 and 2002, the value of the global therapeutic mAb market grew by 37% to US \$5.4 billion. Chimeric mAbs were the undisputed leaders, with 43% growth and US \$3.8 billion in sales, followed by humanized mAbs with more than US \$1.4 billion in sales and growth of 29%. The current global clinical antibody pipeline, which comprises 132 products in development and is dominated by humanized (42%) and fully human (28%) mAbs, is poised to deliver as many as 16 new products between 2004 and 2008. As a result of growth in existing markets for mAb therapeutics, and the opening of new ones, the global market is projected to increase to US \$16.7 billion in 2008.

mAbs approved for marketing

To date, 17 therapeutic mAbs, comprising four different types, have been approved by the US FDA: three murine, five chimeric,




Figure 1 | Phase transition probabilities* for four types of therapeutic mAbs. *Phase transition probabilities were calculated as follows: the number of products that completed a given phase (for example, Phase I) and entered the next (for example, Phase II) was divided by the difference between the number of products that entered the phase and those that were still in the phase at the time of the calculation. Clinical studies for human antibodies initiated during 1994 to 2003. Source: Tufts Center for the Study of Drug Development. USA, US approval; USR, US review.

eight humanized and one human. Nine of the seventeen mAbs have also been approved in the European Union (EU). All of these products were approved in the United States first, though all were approved in the EU within two years of the US approval date. Mean approval times for mAbs are faster in the United States compared with the EU because the FDA has given priority review designation to a majority of the marketing applications. Of the approved products, the best-seller is Johnson & Johnson/Schering-Plough's infliximab (Remicade), with sales of US \$1.6 billion, representing 30.5% of the total market sales in 2002. This product was also the fastest-growing product in 2002, with sales increasing by 84%.

Advancing to the next phase

Although many products might start on the path to approval, not all will complete the convoluted process. In our analysis, probabilities of product advancement from the start of clinical development through US approval were calculated on the basis of current development status of 260 products identified as either murine, chimeric, humanized or human mAbs (FIG. 1). Murine mAbs had the lowest transition probabilities at each phase. The chimeric, humanized and human products had similar Phase I to II and Phase II to III transition probabilities, but the probabilities diverged at the Phase III to US review transition. For the vast majority of mAb products, if the FDA filed the marketing application, then the product ultimately received approval.

Measure of success

Approval success rates are a measure of the likelihood of receiving marketing approval for a product that enters clinical testing. The range of overall approval success for the four types of mAbs was large. To date, the murine products have been least successful (4.5%), whereas the chimeric mAbs have been most successful (26%). The approval success rates for the humanized (18%) and human (14%) mAbs were in the middle of the range. The majority of the humanized and human mAbs are still in clinical development, so the approval success rates might change as the fate of more products is determined. Additional variation in the success rates was observed



'Little Beaver', designed by Frank O'Gehry, from Vitra.com (photographer Hans Hansen)

when the mAbs were categorized by both type and therapeutic category. For example, the success rates for antineoplastic and immunological chimeric mAbs were 29%, whereas humanized antineoplastic and immunological mAbs had a 25% and 17% success rate, respectively.

Waves of the future

Looking toward the future, we anticipate two major approval waves during the next five years. The first will occur between 2004 and 2006, with humanized antibodies comprising the largest number of approvals, whereas the second will occur between 2007 and 2008, and be dominated by human antibody products. Of particular interest, Osidem (ImmunoDesigned Molecules), a combination of a bispecific mAb and macrophage-activated killer cells, and two radiolabelled antibodies might reach the market by 2008. One product produced using a novel engineering approach, Celltech's fragmented antibody CDP-870, is expected to launch in 2006.

A growing market

Although growth will rely on the rise of humanized and human antibodies, chimerics, led by infliximab and rituximab (Rituxan; Genentech), will dominate with a 49% market share in 2008. Humanized antibodies will follow, with sales forecast to reach US \$5.2 billion, or a 31% market share by 2008. In addition, fully human antibodies with 2008 sales of US \$1.9 billion, will capture 11% of the market in 2008.

Two therapeutic categories — oncology and arthritis, immune and inflammatory disorders (AIID) — will likely be the commercial and research focus during the next four years. With the recent approvals of cetuximab (Erbitux; Imclone Systems) and bevacizumab (Avastin; Genentech), oncology will be the leading income earner, with forecast sales of US \$7.2 billion in 2008, representing a 43% market share. Meanwhile, AIID sales will almost quadruple from US \$1.7 billion to \$6.7 billion in 2008, or a 40% market share. In addition, the industry might see approvals in new areas such as the 2005 launch of the humanized antibody natalizumab (Antegren; Biogen IDEC/Elan) for the treatment of multiple sclerosis.

MONOCLONAL ANTIBODIES MARKET | MARKET INDICATORS

has garnered US marketing approval for 13 therapeutic mAbs during the past six years (TABLE 1). During the next five years, the industry has the potential to double the number of approved mAbs, and can anticipate a tripling of the global market for mAb products (see FIGS 2,3). To achieve this result, the industry needs to continue to evolve towards technology integration and market expansion. Success will depend on strategies targeting shorter development times, higher success rates, innovative molecular engineering, robust intellectual property protection and the development of cost-effective manufacturing.

Janice Reichert, Ph.D., is Senior Research Fellow at Tufts Center for the Study of Drug Development, 192 South Street, Suite 550, Boston, Massachusetts 02111, USA. Alex Pavlou, Ph.D., heads Biotechnology Analysis at Datamonitor Healthcare, 108–110 Finchley Road, Charles House, London, NW3 5JJ, UK e-mails: janice.reichert@tufts.edu; apavlou@datamonitor.com

doi:10.1038/nrd1386

 Reichert, J. M. Trends in development and approval times for new therapeutics in the United States. Nature Rev. Drug Discov., 2, 695–702 (2003).

Online links

FURTHER INFORMATION

US Food and Drug Administration: http://www.lda.gov European Agency for the Evaluation of Medicinal Products: http://www.emea.eu.int

Access to this interactive links box is free online.




Figure 2 | New product approval trajectories in terms of technological exposure and sales potential in 2008. Sixteen new antibodies are expected to reach the market over the next four years. Amounts in US \$ millions. Source: Datamonitor. IL, interleukin; MRA, humanized anti-human IL-6 receptor monoclonal antibody.




Figure 3 | Comparison of therapeutic mAb sales trajectories in 2002 (yellow) and 2008 (blue) in terms of technological focus. Despite the undisputed leadership of chimeric mAbs, the contribution of humanized and fully human products to total market size will significantly rise. Source: Datamonitor.

Table 1 | Therapeutic mAbs approved in the United States and European Union

Sponsor company	/ Generic name	US trade name	mAb type	Therapeutic category	US approval date	EU*approval date
Johnson & Johnson	Muromonab-CD3	Orthoclone OKT3	Murine	Immunological‡	19.06.1986	NA
Centocor	Abciximab	ReoPro	Chimeric 🚱	Hemostasis	22.12.1994	YNA *
-Biogen IDEC	Rituximab :	Rituxan	Chimeric	Antineoplastic	26.11.1997.	102:06:1998
Protein Design Labs	Daclizumab	Zenapax	Humanized	Immunological	10.12.1997	26.02.1999
Novártis	Basiliximab	Simulect	-Chimeric 🐉	. Immunological -	12.05.1998.	4.09.10:1998
Medimmune *	Palivizumab	Synagis	Humanized	Anti-Infective	19:06:1998	13.08.1999
Centocor	Infliximab	Remicadé	† Chimerick	: Immunological	24.08:1998	13.08.1999
Genentech	Trastuzumab	Herceptin	Humanized)	Antineoplastic	25:09:1998:	28.08.2000
Wyeth	- Gemtúzumab ozogamicin	Mylotarg	Humanized	Antineoplastic	17.05:2000	INA CLUSS
Millennium/ILEX	Alemtuzumab	Campath	Humanized	Antineoplastic	07.05.2001	06:07:2001
Biogen IDEC	Ibritumomab tiuxetan	Zevalin -	Murine A	Antineoplastic	19.02.2002 年,宋	16.01.2004
Abbott	Adalimumab	Humira .	Human	Immunological	31.12.2002	08:09:2003
Genentech :	Omalizumab :: E	Xólair (:	Humanized	Immunological		ENA TATAL
Cońxa	Tositumomab-I131	BEXXAR	Murine .	Antineoplastic	27.06.2003.	NA .
Genentech.	i Efalizumaba.	Raptiva	Humanized;	Immunological	House 27:10:2003: 2003	L NATA ALTUAR
Imclone Systems	Cetuximab	Erbitux	Chimeric	Antineoplastic	12.02.2004	NA
Genentech,	Bevacizumab	Avastin 1997	Humanized	Antineoplastics	26.02.2004	*NA HAMES

^{*}Approved using EU centralized procedure. *Includes arthritis, immune and inflammatory disorders and prevention/reversal of transplant rejection; NA, not approved. Source: Tufts Center for the Study of Drug Development.



Enablement in Claims to Therapeutic Treatment

Jean Witz tQAS, TC1600



to make and use the invention Specification must teach how

Is the experimentation needed to practice the invention undue or unreasonable?



Therapeutic Treatment

- Inquiry may involve
- How to use the claimed invention
- How to make the claimed invention
- Method claims reciting therapeutic treatment
- Composition or compound claims reciting intended therapeutic use



The amount of guidance or direction knowledge in the state of the art as well as the predictability in the art inversely related to the amount of needed to enable the invention is

However, even in unpredictable arts, a disclosure of every operable species is not required



In re Wands, 858 F.2d 731, 8 USPQ2d 1400 (Fed. Cir. 1988)

Examiner is the fact finder

All evidence must be weighed by the examiner

No per se rules

Case-by-case analysis



- The examiner has the initial burden to question the enablement provided for the either the full scope or a part thereof of the claimed invention establish a reasonable basis to
- contained therein which must be relied There must be a reason to doubt the objective truth of the statements on for enabling support



- possible to support a prima facie References should be supplied if case of lack of enablement, but are not always required
- Specific technical reasons are always required



State of the Art

known in the art at the time of filing Whether or not experimentation is routine depends on what is well-

Enablement analysis is performed presented in the specification based on the state of the art combined with any evidence



State of the Art

technology that is well known at An applicant may omit from the disclosure any routine the time of application



Therapeutic Methods/Uses

unpredictability been resolved by therapeutic method and has this Is there any unpredictability in evidence presented in the the scope of the claimed specification?



In re Gardner, 427 F.2d 786, 166 USPQ 138 (C.C.P.A. 1970)

- Claim to a pharmaceutical composition benzodioxole compounds having comprising 2-aminomethyl-1,3antidepressant activity
- activity, applicants] are claiming in terms of use. It behooves them, therefore, to "In effect, by [claiming therapeutic disclose how to use, as section 112 ordains . . .



In re Gardner, 427 F.2d 786, 166 USPQ 138 (C.C.P.A. 1970)

- Specification lacked the disclosure of
- the recipient of the composition
- the proper dosage
- any working examples
- an animal model



In re Gardner, 427 F.2d 786, 166 USPQ 138 (C.C.P.A. 1970)

- Appellants, relying on an affidavit, argue
- efficacy in a rat model correlated to antidepressant activity in man
- that the proper dosage would have been within the skill of a pharmacologist



Highlights and Guidance

- The lack of direction provided by working examples appeared to be the factors weighed most the inventor and the lack of heavily by the court
- The enablement of compositions reciting activity or intended use must be considered



- myeloblastic leukemia in humans comprising administration of Methods of treating acute naphthacene derivatives
- Pharmaceutical compositions for treatment of acute myeloblastic naphthacene derivatives leukemia comprising



Specification

Claimed derivatives had a close structural relation to daunorubicin and doxorubicin

Two declarations

- Clinical treatment of human patients with acute myeloblastic leukemia
- compounds, one of which was the same Mouse tests on sarcoma tumors and leukemia of eight structurally similar as tested clinically



- The Examiner
- Alleged that there was no utility and therefore no enablement
- Provided no documentary evidence
- The Board
- Affirmed the Examiner
- Provided no documentary evidence



The Court

- noted that neither the solicitor nor the examiner provided support for the assertion regarding "incredible utility"
- held that Board erred by failing to give sufficient weight to the similarity of the remaining claimed derivatives to the allowed derivative
- This similarity combined with the close structural relationship to known antineoplastic agents would have enabled the method/use



Highlights and Guidance

- declaration evidence outweighed amount of direction provided by The state of the prior art, the the Examiner's unsupported the inventor as well as the allegations
- A finding of lack of enablement must be based on evidence



In re Bundy, 642 F.2d 430, 209 USPQ 48 (C.C.P.A. 1981)

- Claims to prostaglandin E analogs
- Specification disclosed
- biological activities of natural PGEs
- therapeutic uses relying on the biological activities
- unexpected increase in analogs' biological activity
- no working examples



In re Bundy, 642 F.2d 430, 209 USPQ 48 (C.C.P.A. 1981)

prostaglandin structure could alter opposed pharmacological effects" potency or induce diametrically enablement citing a reference stating that "small changes in **Examiner found a lack of**



In re Bundy, 642 F.2d 430, 209 USPQ 48 (C.C.P.A. 1981)

- Court held that
- pharmacologic activity was related to PGF, not PGE The evidence of change in
- The discussion of PGE related only to a matter of degree of potency
- compounds without recitation of distinguished due to claims to The result in Gardener was



Highlights and Guidance

compositions that do not recite an intended use need only one Claims to compounds or enabled use

must be sufficiently related to Evidence of unpredictability the claimed invention



Glaxo v. Teva, 2004 WL 1875017 (D. Del. 2004)

- of nausea and vomiting comprising the administration of ondansetron method of treatment for the relief Glaxo patents with claims to a
- As one of the defenses to an action for infringement, Teva asserted lack of enablement of a priority document



Glaxo v. Teva, 2004 WL 1875017 (D. Del. 2004)

working examples in the priority Teva argued the absence of document

The priority document

Identifies ondansetron specifically

Teaches its use as anti-emetic

Provides a dosage range

Provides routes of administration



Glaxo v. Teva, 2004 WL 1875017 (D. Del. 2004)

- **Court finds**
- no requirement in the law for working examples
- priority document clear on its face
- evidence of lack of enablement Teva bore the burden of providing clear and convincing and failed to do so



Highlights and Guidance

Lack of working examples alone is insufficient to support a finding of lack of enablement

may be probative where the evidence indicates unpredictability that may The absence of working examples need to be resolved by exemplary evidence



Rasmussen v. SmithKline, 413 F.3d 1318, 75 USPQ2d 1297 (Fed. Cir. 2005)

- Interference appeal
- Rasmussen lost interference to **SmithKline**
- inhibiting compound, specifically Claims to methods of treating administration of a 5aRprostate cancer by finasteride



Rasmussen v. SmithKline, 413 F.3d 1318, 75 USPQ2d 1297 (Fed. Cir. 2005)

The Board held that Rasmussen's enable the claimed invention in priority document failed to view of

The state of the art

The lack of data to demonstrate the effects of finasteride in treating prostate cancer



Rasmussen v. SmithKline, 413 F.3d 1318, 75 USPQ2d 1297 (Fed. Cir. 2005)

- On appeal, Rasmussen argues that
- showing of efficacy are not relevant to a The Board's findings regarding lack of a finding of lack of enablement, but pertains only to utility
- implausible" that the invention will work The enablement requirement of Section mandates only a showing that it is "not utility and if it does, the requirement 112 does not mandate a showing of for its intended purpose



Rasmussen v. SmithKline, 413 F.3d 1318, 75 USPQ2d 1297 (Fed. Cir. 2005)

- The court disagrees, holding
- may support a rejection under 35 USC 112, 1st paragraph Failure to disclose "how to use"
- unless one with ordinary skill in the art would accept the allegations as "[I]t is proper for the examiner to ask for substantiating evidence obviously correct."



Highlights and Guidance

- was unresolved by the Appellant The unpredictability in the state of the art and the level of skill
- is often the basis for a conclusion that resolves the unpredictability Evidence of unpredictability in the art in the absence of data of lack of enablement



Impax v. Aventis, 496 F.Supp.2d 428 (D. Del. 2007)

- Claims to method of treating ALS by administering riluzole
- Impax asserted invalidity based on prior art anticipation of **Aventis patent**
- Aventis argued asserted prior art was not enabling



Impax v. Aventis, 392 F.Supp.2d 428 (D. Del. 2007)

- Aventis asserted that the patent
- compounds and numerous diseases, discloses thousands of formula I yielding thousands of possible combinations
- provides no direction or guidance to arrive at the claimed invention of using riluzole to treat ALS
- examples of the claimed invention does not disclose any working



Impax v. Aventis, 392 F.Supp.2d 428 (D. Del. 2007)

- Impax asserted that the patent
- includes riluzole as a formula I compound
- suggests that formula I compounds may be used to treat ALS
- provides some dosage information



Impax v. Aventis, 392 F.Supp.2d 428 (D. Del. 2007)

contained in the patent to suggest that Impax directs the Court to information undue experimentation would not be required

In human therapy, the compounds according to the invention are especially useful in the treatment and prevention of convulsive phenomena, schizophrenic disorders, and in particular the deficiency forms of schizophrenia, sleep disorders, phenomena linked to cerebral ischaemia and also neurological conditions in which glutamate may be implicated, such as Alzehimer's disease, Huntington's chorea, amyotrophic lateral sclerosis and olivopontocerebellar atrophy



Impax v. Aventis, 392 F.Supp.2d 428 (D. Del. 2007)

The District Court finds

- different diseases, and there is nothing in the patent which would lead one to recognize that any specific compound, let alone riluzole, "the compounds of the claimed invention are would be used to treat any specific disease, associated with the treatment of at least 8 let alone ALS."
 - that the mere mention of riluzole was insufficient to put one skilled in the art in the possession of the claimed invention as is required to support a conclusion of enablement



Highlights and Guidance

compounds to be used, yielding large Specification detailing extensive lists evidence of unpredictability exists in absence of working examples and if claim to specific combination in the may suggest lack of enablement of numbers of possible combinations of conditions to be treated and the prior art



Laboratories, Inc., 2007 WL 3151692 (Fed. Cir. 2007) Pharmaceutical Resources v. Roxane

- affirming the District Court finding Non-precedential Fed. Cir. opinion that Par's patents were invalid for lack of enablement
- composition of megestrol acetate, choices of specific alcohols and a Claims to oral pharmaceutical surfactant



Laboratories, Inc., 2007 WL 3151692 Pharmaceutical Resources v. Roxane (Fed. Cir. 2007)

- Claim language did not limit type or amount of surfactant
- Specification stated that invention was not limited to particular surfactants
- Par asserted that broadest reasonable interpretation of claim did not limit type or amount of surfactant



Laboratories, Inc., 2007 WL 3151692 Pharmaceutical Resources v. Roxane (Fed. Cir. 2007)

- Par stressed unpredictability in formulation based on type and amount of surfactant during prosecution of patents
- with surfactants during previous unpredictability of formulation trial with another litigant Par's expert testified to



Laboratories, Inc., 2007 WL 3151692 Pharmaceutical Resources v. Roxane (Fed. Cir. 2007)

- evidence of unpredictability provided The court held the claims lacked enablement based, in part, on previously by Par
- The court also considered the breadth working examples and unsupported of the claims, the presence of conclusions in declarations



Highlights and Guidance

enablement for at least a portion conclusion of nonobviousness Evidence of unpredictability may be then appropriate to support a finding of lack of of the scope of the claim presented to support a





therapeutic treatment claims determining breadth of the begins with the claims by **Enablement analysis of** claims with regard to

The condition to be treated

The compound/composition administered



Review

continues with determination of unpredictability within the state therapeutic treatment claims of the art with regard to **Enablement analysis of** the presence of any

The condition to be treated

The compound/composition administered



Review

- **Enablement analysis of therapeutic** treatment claims finishes with the specification by evaluation of
- The presence or absence of working examples
- The evaluation of any other evidence of record, e.g. declarations



Review

- Evidence of unpredictability or predictability may occur in the
- Etiology of the condition/disease
- Number/type of other accepted treatments
- The presence or absence of artrecognized animal models
- Manner of formulation and/or delivery



Highlights and Guidance

- and must provide the evidence The Examiner is the fact finder
- The Examiner must weigh the evidence and provide the rationale
- No per se rules!



Highlights and Guidance

- Consider claim construction
- Consider the evidence
- No per se rules!



Thank You!

jean.witz@uspto.gov

571-272-0927



Enablement in Claims to Therapeutic Treatment

Jean Witz tQAS, TC1600



- to make and use the invention Specification must teach how
- Is the experimentation needed to practice the invention undue or unreasonable?



Therapeutic Treatment

- Inquiry may involve
- How to use the claimed invention
- How to make the claimed invention
- Method claims reciting therapeutic treatment
- reciting intended therapeutic use Composition or compound claims



The amount of guidance or direction knowledge in the state of the art as well as the predictability in the art inversely related to the amount of needed to enable the invention is

However, even in unpredictable arts, a disclosure of every operable species is not required



In re Wands, 858 F.2d 731, 8 USPQ2d 1400 (Fed. Cir. 1988)

Examiner is the fact finder

All evidence must be weighed by the examiner

No per se rules

Case-by-case analysis



- The examiner has the initial burden to question the enablement provided for the either the full scope or a part thereof of the claimed invention establish a reasonable basis to
- contained therein which must be relied There must be a reason to doubt the objective truth of the statements on for enabling support



- possible to support a prima facie References should be supplied if case of lack of enablement, but are not always required
- Specific technical reasons are always required



State of the Art

- known in the art at the time of filing Whether or not experimentation is routine depends on what is well-
- Enablement analysis is performed presented in the specification combined with any evidence based on the state of the art



State of the Art

technology that is well known at the time of application An applicant may omit from the disclosure any routine



Therapeutic Methods/Uses

unpredictability been resolved by therapeutic method and has this Is there any unpredictability in evidence presented in the the scope of the claimed specification?



In re Gardner, 427 F.2d 786, 166 USPQ 138 (C.C.P.A. 1970)

- Claim to a pharmaceutical composition benzodioxole compounds having comprising 2-aminomethyl-1,3antidepressant activity
- activity, applicants] are claiming in terms of use. It behooves them, therefore, to "In effect, by [claiming therapeutic disclose how to use, as section 112 ordains . . .



In re Gardner, 427 F.2d 786, 166 USPQ 138 (C.C.P.A. 1970)

Specification lacked the disclosure of the recipient of the composition

the proper dosage

any working examples

an animal model



In re Gardner, 427 F.2d 786, 166 USPQ 138 (C.C.P.A. 1970)

- Appellants, relying on an affidavit, argue
- efficacy in a rat model correlated to antidepressant activity in man
- that the proper dosage would have been within the skill of a pharmacologist



- The lack of direction provided by working examples appeared to be the factors weighed most the inventor and the lack of heavily by the court
- The enablement of compositions reciting activity or intended use must be considered



- myeloblastic leukemia in humans comprising administration of Methods of treating acute naphthacene derivatives
- Pharmaceutical compositions for treatment of acute myeloblastic naphthacene derivatives leukemia comprising



Specification

Claimed derivatives had a close structural relation to daunorubicin and doxorubicin

Two declarations

- Clinical treatment of human patients with acute myeloblastic leukemia
- compounds, one of which was the same leukemia of eight structurally similar Mouse tests on sarcoma tumors and as tested clinically



- The Examiner
- Alleged that there was no utility and therefore no enablement
- Provided no documentary evidence
- The Board
- Affirmed the Examiner
- Provided no documentary evidence



The Court

- noted that neither the solicitor nor the examiner provided support for the assertion regarding "incredible utility"
- held that Board erred by failing to give sufficient weight to the similarity of the remaining claimed derivatives to the allowed derivative
- This similarity combined with the close structural relationship to known antineoplastic agents would have enabled the method/use



declaration evidence outweighed amount of direction provided by The state of the prior art, the the Examiner's unsupported the inventor as well as the allegations

A finding of lack of enablement must be based on evidence



In re Bundy, 642 F.2d 430, 209 USPQ 48 (C.C.P.A. 1981)

Claims to prostaglandin E analogs

Specification disclosed

biological activities of natural PGEs

therapeutic uses relying on the biological activities unexpected increase in analogs' biological activity

no working examples



In re Bundy, 642 F.2d 430, 209 USPQ 48 (C.C.P.A. 1981)

prostaglandin structure could alter opposed pharmacological effects" potency or induce diametrically enablement citing a reference stating that "small changes in **Examiner found a lack of**



In re Bundy, 642 F.2d 430, 209 USPQ 48 (C.C.P.A. 1981)

- Court held that
- pharmacologic activity was related to PGF, not PGE The evidence of change in
- The discussion of PGE related only to a matter of degree of potency
- compounds without recitation of distinguished due to claims to The result in Gardener was



compositions that do not recite an intended use need only one Claims to compounds or enabled use

must be sufficiently related to Evidence of unpredictability the claimed invention



Glaxo v. Teva, 2004 WL 1875017 (D. Del. 2004)

- of nausea and vomiting comprising the administration of ondansetron method of treatment for the relief Glaxo patents with claims to a
- As one of the defenses to an action for infringement, Teva asserted lack of enablement of a priority document



Glaxo v. Teva, 2004 WL 1875017 (D. Del. 2004)

working examples in the priority Teva argued the absence of document

The priority document

Identifies ondansetron specifically

Teaches its use as anti-emetic

Provides a dosage range

Provides routes of administration



Glaxo v. Teva, 2004 WL 1875017 (D. Del. 2004)

- Court finds
- no requirement in the law for working examples
- priority document clear on its race
- evidence of lack of enablement providing clear and convincing Teva bore the burden of and failed to do so



- Lack of working examples alone is insufficient to support a finding of lack of enablement
- may be probative where the evidence indicates unpredictability that may The absence of working examples need to be resolved by exemplary evidence



Rasmussen v. SmithKline, 413 F.3d 1318, 75 USPQ2d 1297 (Fed. Cir. 2005)

- Interference appeal
- Rasmussen lost interference to **SmithKline**
- inhibiting compound, specifically Claims to methods of treating administration of a 5aRprostate cancer by finasteride



Rasmussen v. SmithKline, 413 F.3d 1318, 75 USPQ2d 1297 (Fed. Cir. 2005)

The Board held that Rasmussen's enable the claimed invention in priority document failed to view of

The state of the art

The lack of data to demonstrate the effects of finasteride in treating prostate cancer



Rasmussen v. SmithKline, 413 F.3d 1318, 75 USPQ2d 1297 (Fed. Cir. 2005)

- On appeal, Rasmussen argues that
- showing of efficacy are not relevant to a The Board's findings regarding lack of a finding of lack of enablement, but pertains only to utility
- implausible" that the invention will work The enablement requirement of Section mandates only a showing that it is "not utility and if it does, the requirement 112 does not mandate a showing of for its intended purpose



Rasmussen v. SmithKline, 413 F.3d 1318, 75 USPQ2d 1297 (Fed. Cir. 2005)

- The court disagrees, holding
- may support a rejection under 35 USC 112, 1st paragraph Failure to disclose "how to use"
- unless one with ordinary skill in the art would accept the allegations as "[I]t is proper for the examiner to ask for substantiating evidence obviously correct."



- was unresolved by the Appellant The unpredictability in the state of the art and the level of skill
- is often the basis for a conclusion that resolves the unpredictability Evidence of unpredictability in the art in the absence of data of lack of enablement



- Claims to method of treating ALS by administering riluzole
- Impax asserted invalidity based on prior art anticipation of **Aventis** patent
- Aventis argued asserted prior art was not enabling



- Aventis asserted that the patent
- compounds and numerous diseases, discloses thousands of formula I yielding thousands of possible combinations
- provides no direction or guidance to arrive at the claimed invention of using riluzole to treat ALS
 - examples of the claimed invention does not disclose any working



- Impax asserted that the patent
- includes riluzole as a formula I punodwoo
- suggests that formula I compounds may be used to treat ALS
- provides some dosage information



contained in the patent to suggest that undue experimentation would not be Impax directs the Court to information required

In human therapy, the compounds according to the invention are especially useful in the treatment and prevention of convulsive phenomena, schizophrenic disorders, and in particular the deficiency forms of schizophrenia, sleep disorders, phenomena linked to cerebral ischaemia and also neurological conditions in which glutamate may be implicated, such as Alzehimer's disease, Huntington's chorea, amyotrophic lateral scierosis olivopontocerebellar atrophy



The District Court finds

that any specific compound, let alone riluzole, different diseases, and there is nothing in the patent which would lead one to recognize "the compounds of the claimed invention are would be used to treat any specific disease, let alone ALS." associated with the treatment of at least 8

insufficient to put one skilled in the art in the possession of the claimed invention as is that the mere mention of riluzole was required to support a conclusion of enablement



compounds to be used, yielding large Specification detailing extensive lists evidence of unpredictability exists in absence of working examples and if claim to specific combination in the may suggest lack of enablement of numbers of possible combinations of conditions to be treated and the prior art



Laboratories, Inc., 2007 WL 3151692 (Fed. Cir. 2007) Pharmaceutical Resources v. Roxane

- affirming the District Court finding Non-precedential Fed. Cir. opinion that Par's patents were invalid for lack of enablement
- composition of megestrol acetate, choices of specific alcohols and a Claims to oral pharmaceutical surfactant



Laboratories, Inc., 2007 WL 3151692 Pharmaceutical Resources v. Roxane (Fed. Cir. 2007)

- Claim language did not limit type or amount of surfactant
- Specification stated that invention was not limited to particular surfactants
- Par asserted that broadest reasonable interpretation of claim did not limit type or amount of surfactant



Laboratories, Inc., 2007 WL 3151692 (Fed. Cir. 2007) Pharmaceutical Resources v. Roxane

- Par stressed unpredictability in formulation based on type and amount of surfactant during prosecution of patents
- with surfactants during previous unpredictability of formulation trial with another litigant Par's expert testified to



Laboratories, Inc., 2007 WL 3151692 Pharmaceutical Resources v. Roxane (Fed. Cir. 2007)

- evidence of unpredictability provided The court held the claims lacked enablement based, in part, on previously by Par
- The court also considered the breadth working examples and unsupported of the claims, the presence of conclusions in declarations



enablement for at least a portion conclusion of nonobviousness Evidence of unpredictability may be then appropriate to support a finding of lack of of the scope of the claim presented to support a



therapeutic treatment claims determining breadth of the begins with the claims by **Enablement analysis of** claims with regard to

The condition to be treated

The compound/composition administered



continues with determination of unpredictability within the state therapeutic treatment claims of the art with regard to **Enablement analysis of** the presence of any

The condition to be treated

The compound/composition administered



- **Enablement analysis of therapeutic** treatment claims finishes with the specification by evaluation of
- The presence or absence of working examples
- The evaluation of any other evidence of record, e.g. declarations



- Evidence of unpredictability or predictability may occur in the
- Etiology of the condition/disease
- Number/type of other accepted treatments
- The presence or absence of artrecognized animal models
- Manner of formulation and/or delivery



and must provide the evidence The Examiner is the fact finder

The Examiner must weigh the evidence and provide the rationale

No per se rules!



- Consider claim construction
- Consider the evidence
- No per se rules!



Thank You!

i jean.witz@uspto.gov

571-272-0927